

Final Report

Bacterial Reverse Mutation Screening Assay using *Salmonella typhimurium*

Test Article: GelRed (lot # BT-17-119-FM)

Date Initiated: February 21, 2006

Date Completed: March 16, 2006

Protocol: A70S-2006

Litron's Study Director: Daniel J. Roberts, B.S.

Testing Facility: Litron Laboratories
200 Canal View Boulevard
Suite 106
Rochester, New York 14623

Sponsor: Biotium, Inc.
3423 Investment Boulevard
Suite 8
Hayward, California 94545

Sponsor Representative: Dr. Xing Xin
Research and Development

Table of Contents

1.	List of Data Tables and Figures	3
2.	Objective	4
3.	Test Article and Vehicle Description	4
4.	Summary	4
5.	Materials and Methods	5
5.1	Test System Description	5
5.2	Test System Justification.....	5
5.3	Source and Storage of Test System.....	5
5.4	Identification of Test System	5
5.5	Preparation of Overnight Cultures	5
5.6	Control of Bias.....	5
5.7	Metabolic Activation.....	6
5.8	Tester Strain Media	6
5.9	Definitive Assay	6
5.10	Repeat Assay.....	7
5.11	Counting	8
5.11	Criteria for a Valid Assay.....	8
5.12	Statistical Analysis of the Data	8
5.13	Determining a Positive Response	9
6.	Historical Control Data.....	9
7.	Legend	9
8.	Discussion of Results	10
9.	Conclusions	10
10.	Records Maintained	10
11.	References	10
	Data Tables and Figures	11
	Approved Protocol	Attached

1. List of Data Tables and Figures

Table 1: Control Data (Definitive Assay).....	11
Table 2: Control Data (Repeat Assay)	12
Figure and Table 3: Strain TA98 (Definitive and Repeat Assays)	13
Figure and Table 4: Strain TA1537 (Definitive Assay).....	14
Figure and Table 5: Strain TA98 (Definitive Assay – Incomplete Data Set)	15

2. Objective

This Bacterial Reverse Mutation Screening Assay was performed to evaluate the ability of this test article to induce a mutagenic response in two strains of *Salmonella typhimurium* (TA98 and TA1537).

3. Test Article and Vehicle Description

Test article characteristics: Test article was received as a dark purple, almost black solution in a clear vial labeled: "Cat: 41000, Lot: BT-17-119-FM, 0.9 ml, GelRed, 10mg/ml in DMSO, Biotium, Inc., For Research Use Only." The protocol indicated that this test article expires 2/15/2007.

Storage conditions: The test article was stored at -24 °C to -21 °C, protected from light.

Vehicle (and lot number): Dimethyl sulfoxide (Lot: 05754KD, Exp: 9/07).

Justification for vehicle choice: Sponsor indicated vehicle to be utilized.

Description of test article when mixed with vehicle: A dark red solution.

4. Summary

The results of this Bacterial Reverse Mutation Screening Assay indicate that under the experimental conditions, this test article was mutagenic for tester strain TA98 with S9 metabolic activation.

Protocol A70S-2006 is based on Organisation for Economic Co-operation and Development (OECD) and International Conference on Harmonisation (ICH) testing guidelines.

5. Materials and Methods

5.1. Test System Description

The Salmonella strains used were histidine-dependent. Revertants were identified as colonies that grew in low levels of histidine. Frameshift substitution defects were represented to identify mutagens. Additional genetic markers enhanced sensitivity of the strains to certain types of mutagens.

The DNA repair mutation (*uvrB*) eliminates excision repair, a repair pathway for DNA damage from UV light and certain chemical mutagens. The *uvrB* mutation, present in both strains, was indicated by sensitivity to UV light. The *rfa* mutation changes the properties of the bacterial cell wall, increasing permeability of cells to certain types of chemicals. The *rfa* mutation, present in both strains was indicated by sensitivity to crystal violet.

The R factor plasmid (pKM101) present in strain TA98 makes it more responsive to a variety of mutagens. The plasmid carries an ampicillin resistance gene; therefore ampicillin resistance indicated that the strain retain the plasmid.

CHARACTERISTICS OF TESTER STRAINS						
Tester Strain	Gene Affected	DNA Repair	LPS	Biotin Requirement	Plasmids	Mutational Event
TA98	<i>hisD</i>	<i>uvrB</i>	<i>rfa</i>	bio-	pKM101	frameshift
TA1537	<i>hisC</i>	<i>uvrB</i>	<i>rfa</i>	bio-	-	frameshift

5.2. Test System Justification

The two strains of bacteria used in this assay are among those recommended by OECD 471 for use in the Ames test. These two strains of *S. typhimurium* have been shown to be reliably and reproducibly responsive between laboratories.

5.3. Source and Storage of Test System

The Salmonella strains used in this study were obtained from Xenometrix and Molecular Toxicology, Inc. Cells are maintained as frozen stocks (-85 °C ± 4 °C).

5.4. Identification of Test System

Strains TA98 and TA1537 were identified by having certain characteristics (see above). The strains also yielded spontaneous revertant colony plate counts within the frequency ranges stated in the historical control data.

5.5. Preparation of Overnight Cultures

Frozen stock cultures (stored at -85 °C ± 4 °C) were grown overnight at 37 ± 2 °C, with shaking, in nutrient broth until a cell density of 1 x 10⁹ to 2 x 10⁹ cells/ml is obtained (determined by optical density). Cells were refrigerated until use and maintained at room temperature during the test.

5.6. Control of Bias

In order to control bias on the day of test system treatment, all test article doses, as well as controls, were plated against cells obtained from a single flask for each strain.

5.7. Metabolic Activation

5.7.1. S9 Fraction

Aroclor™ 1254-induced male Sprague Dawley rat liver S9 (500 mg/kg i.p.), was purchased from a commercial supplier (Molecular Toxicology, Inc. Boone, NC). Lot 1938 was used, which contains 40.8 mg/ml protein. This lot has demonstrated the ability to activate ethidium bromide, cyclophosphamide, benzo(a)pyrene, and 2-aminoanthracene into mutagenic intermediates, and has demonstrated P450 substrate activation in the ethoxyresorufin-O-deethylase, pentoxyresorufin-O-dealkylase, benzylresorufin-O-dealkylase and methoxyresorufin-O-dealkylase assays.

5.7.2. S9/Cofactor Mix

The S9/Cofactor Mix was prepared immediately before the test and contained: 10% S9, magnesium chloride, potassium chloride, D-glucose-6-phosphate, and nicotinamide adenine dinucleotide phosphate, in a sodium phosphate buffer. It was kept on ice during the experiment.

5.7.3. Buffer

When S9 mix was not used in the test, phosphate buffered saline (PBS) was used in its place.

5.8. Tester Strain Media

5.8.1. Nutrient Broth

The broth used for the overnight cultures consisted of 2.5% Oxoid Nutrient Broth #2.

5.8.2. Vogel-Bonner Plates

Minimal glucose agar plates (1.5% agar supplemented with 2.0% glucose and 2.0% Vogel-Bonner buffer) were purchased from a commercial supplier (Moltox, Boone, NC).

5.8.3. Top Agar

Top agar was prepared with 0.6% agar and 0.6% NaCl supplemented with histidine (0.5 mM) and biotin (0.5 mM). For the assay, 2.0 ml supplemented top agar was used.

5.9. Definitive Assay

Concentrations of test article prepared: 0, 1, 2.5, 5, 10, 25, 50, 75, 100, 250 and 500 µg/ml.

Volume of each concentration plated: 0.1 ml per plate.

Doses tested: 0, 0.1, 0.25, 0.5, 1, 2.5, 5, 7.5, 10, 25 and 50 µg/plate.

Justification for doses tested: Sponsor requested these concentrations be tested.

Number of plates per dose: Duplicate plates were used for each dose.

Microbiological contamination: An aliquot of the top concentration was plated to test for microbiological contamination. None was evident.

The following was added to each sterile culture tube containing 2.0 ml top agar: 0.1 ml of overnight cell culture (TA98 or TA1537), 0.1 ml of each test article concentration or control chemical, and either 0.5 ml of S9/Cofactor mix or 0.5 ml of phosphate buffered saline.

An insufficient volume of the TA98 cell culture was available to plate the last five plates of this test article with S9 metabolic activation (one plate of 10 $\mu\text{g}/\text{plate}$ and duplicates of 25 and 50 $\mu\text{g}/\text{plate}$). In order to obtain a complete data set to evaluate the mutagenicity of this test article, all concentrations along with concurrent controls for strain TA98 with S9 metabolic activation only, were repeated. See section 5.10.

The contents of each tube were vortexed, poured onto Vogel-Bonner media plates, and evenly distributed. The agar on the test plates was allowed to harden. The plates were inverted and incubated at $37\text{ }^{\circ}\text{C} \pm 2\text{ }^{\circ}\text{C}$ for 48 hours \pm 4 hours prior to scoring.

Was test article precipitation observed? If yes, at what doses? No test article precipitation was observed.

Was toxicity observed? If yes, at what doses? No toxicity was observed.

5.10. Repeat Assay

Concentrations of test article prepared: 0, 1, 2.5, 5, 10, 25, 50, 75, 100, 250 and 500 $\mu\text{g}/\text{ml}$.

Volume of each concentration plated: 0.1 ml per plate.

Doses tested: 0, 0.1, 0.25, 0.5, 1, 2.5, 5, 7.5, 10, 25 and 50 $\mu\text{g}/\text{plate}$.

Justification for doses tested: Sponsor requested these concentrations be tested.

Number of plates per dose: Duplicate plates were used for each dose.

Microbiological contamination: An aliquot of the top concentration was plated to test for microbiological contamination. None was evident.

The following was added to each sterile culture tube containing 2.0 ml top agar: 0.1 ml of overnight cell culture (TA98), 0.1 ml of each test article concentration or control chemical, and 0.5 ml of S9/Cofactor mix.

The contents of each tube were vortexed, poured onto Vogel-Bonner media plates, and evenly distributed. The agar on the test plates was allowed to harden. The plates were inverted and incubated at $37\text{ }^{\circ}\text{C} \pm 2\text{ }^{\circ}\text{C}$ for 48 hours \pm 4 hours prior to scoring.

Was test article precipitation observed? If yes, at what doses? No test article precipitation was observed.

Was toxicity observed? If yes, at what doses? No toxicity was observed.

5.11. Counting

5.11.1. Automatic Colony Counting

A New Brunswick Biotran III automatic colony counter was used for counting revertant colonies. The control chemical plates were counted before the test article plates for each strain to ensure that the strain was functioning properly. Before counting, each plate was scanned for contamination, test article precipitation, toxicity and any foreign material. Toxicity is suggested by the absence of a confluent bacterial lawn, the presence of pinpoint colonies, and/or a substantial decrease or lack of revertant colonies. Each plate was counted 3 times on the automatic colony counter, rotating the plate one third each time. The median count was recorded.

5.11.2. Hand Counting

Hand counting was not required.

5.12. Criteria for a Valid Assay

The study will be considered valid if the following criteria are met.

- Both tester strains demonstrated the presence of the *uvrB* mutation by exhibiting sensitivity to UV light.
- Both tester strains demonstrated the presence of the *rfa* wall mutation by exhibiting sensitivity to crystal violet.
- Tester strain TA98 demonstrated the presence of the pKM101 plasmid by exhibiting resistance to ampicillin.
- Each tester strain demonstrated a characteristic number of spontaneous revertant colonies. A “characteristic number” is defined as the average number of colonies across plates being within the historical range, or within the published historical range.
- Each tester strain exhibited at least a three-fold increase in average mutagen-induced revertant colonies when plated with positive control chemicals.

5.13. Statistical Analysis of the Data

Only when an assay is valid, and only when any test article treatment group demonstrates an increase in average number of revertant colonies relative to the negative control, will data be subjected to statistical analyses.

The average of each set of duplicate plates was determined. JMP software’s regression analysis (v5) was used to determine if a dose-related increase occurred ($p < 0.025$). Statistically significant dose-related increases were observed for strains TA98 ($p < 0.0001$) and TA1537 ($p = 0.0041$) with S9 metabolic activation.

5.14. Determining a Positive Response

The test article will be considered positive if the assay is valid, and if the following conditions are met, taking into account biological relevance:

- One test article dose exceeds three times the background average (two times for strain TA1537) either with or without metabolic activation, or there is a dose-related increase over the range tested ($p < 0.025$).
- If the background average is below six colonies, the average number of revertants for the test article must exceed 20 colonies/plate.

A positive result indicates that the test article induces mutations in *Salmonella typhimurium* cells.

A test article for which the results do not meet the above criteria will be considered non-mutagenic in this test. Negative results indicate that, under the test conditions, the test article does not produce mutations in *Salmonella typhimurium* cells.

Although most experiments will give clearly positive or negative results, in rare cases the data set will preclude making a definite judgment about the activity of the test article. Results may remain equivocal or questionable regardless of the number of times the experiment is repeated.

6. Historical Control Data

Average Number of Colonies per Plate ± Standard Deviation (Range Indicated Below)		
Agent	<i>Salmonella typhimurium</i> strains	
	TA98	TA1537
Top Agar + His	11 ± 3 (7 - 16)	6 ± 2 (2 - 10)
DMSO	11 ± 5 (4 - 22)	5 ± 2 (1 - 9)
Positive Control (µg/plate)	2NF (1)	9AA (50)
	224 ± 65 (132 - 313)	93 ± 49 (40 - 225)
Top Agar + 10% S9	19 ± 5 (9 - 28)	4 ± 2 (1 - 7)
DMSO + 10% S9	18 ± 4 (13 - 26)	5 ± 2 (1 - 8)
Positive Control + 10% S9 (µg/plate)	BP (3)	2AAn (5)
	281 ± 25 (229 - 342)	144 ± 21 (96 - 171)

Historical data determined from six Ames tests, February through March 2006.

7. Legend

His:	histidine	
S9:	S9 metabolic activation	
DMSO:	dimethyl sulfoxide	CAS#: 67-68-5
2NF:	2-nitrofluorene	CAS#: 607-57-8
NaN ₃ :	sodium azide	CAS#: 26628-22-8
BP:	benzo(a)pyrene	CAS#: 50-32-8
2AAn:	2-aminoanthracene	CAS#: 613-13-8

8. Discussion of Results

Although a statistically significant dose-related increase was observed with strain TA1537, with S9 metabolic activation, the increase is not considered biologically relevant for the following reasons: the highest average number of revertant colonies (9) was within Litron's historical range for this strain (7 – 16); and, the background average for the vehicle control was below six colonies, while no test article concentration had an average number of revertants above 20 (see Section 5.14).

A statistically significant increase was observed for strain TA98 with S9 metabolic activation.

9. Conclusions

The results of this assay indicate that under the experimental conditions, this test article was mutagenic for tester strain TA98 with S9 metabolic activation.

10. Records Maintained

All records regarding the study, including correspondence between the sponsor and Litron, the protocol, amendments to the protocol, data sheets, environmental and equipment information, training records, historical data, a copy of the final report, and all other raw data and applicable information, will be maintained at Litron for five years following completion of the study. Electronic copies of records will be stored off-site (315 Root Rd., Brockport, NY 14420) in addition to storage at Litron Laboratories.

11. References

- Ames, B., J. McCann, and E. Yamasaki (1975) Methods for detecting carcinogens and mutagens with the Salmonella/mammalian-microsome mutagenicity test. *Mutation Res.* 31, 347-364.
- Gatehouse, D., S. Haworth, T. Cebula, E. Gocke, L. Kier, T. Matsushima, C. Melcion, T. Nohmi, S. Venitt, and E. Zeiger (1994) Recommendations for the performance of bacterial mutation assays. *Mutation Res.* 312, 217-233.
- Mahon, G., M. Green, B. Middleton, I. Mitchell, W. Robinson, and D. Tweats (1989) Analysis of Data from Microbial Colony Assays. In: UKEMS Sub-Committee on Guidelines for Mutagenicity Testing Part II. Statistical Evaluation of Mutagenicity Test Data. Ed. Kirkland, D. J., Cambridge Press, pp. 28-65.
- Majeska, J. and J. Mayo (2000) Poster presented at the Thirty-First Annual Meeting of the Environmental Mutagen Society.
- Maron, D. and B. Ames (1983) Revised methods for the Salmonella mutagenicity test. *Mutation Res.* 113, 173-215.
- Maron, D., J. Katzenellenbogen, and B. Ames (1981) Compatibility of organic vehicles with the Salmonella/microsome test. *Mutation Res.* 88, 343-350. Maron, D., J. Katzenellenbogen, and B. Ames (1981) *Mutation Res.* 88, 343-350.
- McKee, R., J. Tometsko, and A. Tometsko (1979) Chemicals which revert all commonly used *Salmonella typhimurium* tester strains. *Mutation Res.* 67, 183-187.
- Mortelmans K. and Ziegler E. (2000) *Mutation Res.* 455, 29-60.
- Wilcox, P., A. Naidoo, D. Wedd, and D. Gatehouse (1990) Comparison of Salmonella typhimurium TA102 with Escherichia coli WP2 tester strains. *Mutagenesis* 5, 285-291.
- International Conference on Harmonisation (ICH) Tripartite Guidelines: Guidance on Specific Aspects of Regulatory Genotoxicity Tests for Pharmaceuticals. S2A, adopted July 19, 1995.
- International Conference on Harmonisation (ICH) Tripartite Guidelines: Genotoxicity: A Standard Battery for Genotoxicity Testing of Pharmaceuticals. S2B, adopted July 16, 1997.
- Organisation for Economic Cooperation and Development (OECD) Section 4 of the OECD Guidelines for the Testing of Chemicals: Bacteria Reverse Mutation Test, Guideline 471, adopted July 21, 1997.

Table 1: Control Data (Definitive Assay)

Date plated: March 1, 2006

Date counted: March 3, 2006

Raw Data (Number of Colonies per Plate)

	<i>Salmonella typhimurium</i> Strains			
	TA98		TA1537	
Top Agar + His	8	10	C	5
DMSO	10	11	2	1
Positive Control	2NF (1 µg/plate)		9AA (50 µg/plate)	
	172	187	203	190
Top Agar + S9	15	15	6	8
DMSO +S9	13	15	4	7
Positive Control + S9	BP (3 µg/plate)		2AAAn (5 µg/plate)	
	242	250	157	152

Average Number of Colonies per Plate

	<i>Salmonella typhimurium</i> Strains	
	TA98	TA1537
Top Agar + His	9	5
DMSO	11	2
Positive Control	2NF (1 µg/plate)	9AA (50 µg/plate)
	180	197
Top Agar + S9	15	7
DMSO +S9	14	6
Positive Control + S9	BP (3 µg/plate)	2AAAn (5 µg/plate)
	246	155

C = Contamination.

Table 2: Control Data (Repeat Assay)

Date plated: March 8, 2006

Date counted: March 10, 2006

Raw Data (Number of Colonies per Plate)

	<i>Salmonella typhimurium</i> Strain	
	TA98	
Top Agar + S9	20	18
DMSO +S9	30	23
Positive Control + S9	BP (3 µg/plate)	
	246	303



Average Number of Colonies per Plate

	<i>Salmonella typhimurium</i> Strain	
	TA98	
Top Agar + S9	19	
DMSO +S9	27	
Positive Control + S9	BP (3 µg/plate)	
	275	

Figure and Table 3: Strain TA98 (Definitive and Repeat Assays)

Date plated: March 1, 2006 and †March 8, 2006

Date counted: March 3, 2006 and †March 10, 2006





Dose (µg/plate)	Number of Colonies	Averages
0	8 9	9
0.1	5 12	9
0.25	14 7	11
0.5	7 9	8
1	5 7	6
2.5	8 12	10
5	9 11	10
7.5	4 4	4
10	15 10	13
25	11 7	9
50	7 9	8

Dose (µg/plate)	Number of Colonies	Averages
0 + S9	24 17	21
0.1 + S9	16 20	18
0.25 + S9	30 20	25
0.5 + S9	24 24	24
1 + S9	20 26	23
2.5 + S9	18 24	21
5 + S9	17 22	20
7.5 + S9	27 27	27
10 + S9	22 28	25
25 + S9	42 37	40
50 + S9	59 48	54

Figure and Table 4: Strain TA1537 (Definitive Assay)

Date plated: March 1, 2006

Date counted: March 3, 2006




Dose (µg/plate)	Number of Colonies		Averages
0	3	1	2
0.1	3	3	3
0.25	3	5	4
0.5	1	6	4
1	2	4	3
2.5	4	4	4
5	1	2	2
7.5	2	4	3
10	9	3	6
25	2	7	5
50	6	5	6

Dose (µg/plate)	Number of Colonies		Averages
0 + S9	5	5	5
0.1 + S9	1	4	3
0.25 + S9	3	7	5
0.5 + S9	5	2	4
1 + S9	5	6	6
2.5 + S9	7	5	6
5 + S9	7	1	4
7.5 + S9	7	6	7
10 + S9	6	5	6
25 + S9	6	10	8
50 + S9	7	10	9

Figure and Table 5: Strain TA98 (Definitive Assay – Incomplete Data Set)

Date plated: March 1, 2006

Date counted: March 3, 2006



Dose (µg/plate)	Number of Colonies	Averages
0 + S9	15 16	16
0.1 + S9	13 20	17
0.25 + S9	9 10	10
0.5 + S9	12 15	14
1 + S9	14 10	12
2.5 + S9	15 16	16
5 + S9	15 19	17
7.5 + S9	18 13	16
10 + S9	25	25